



Perspective

COVID toes: Where do we stand with the current evidence?

Marie Baeck*, Anne Herman

Department of Dermatology, Cliniques universitaires Saint-Luc, Université Catholique de Louvain (UCLouvain), Avenue Hippocrate 10, 1200 Brussels, Belgium



ARTICLE INFO

Article history:

Received 11 September 2020

Received in revised form 5 October 2020

Accepted 8 October 2020

Keywords:

COVID-19

SARS-CoV-2

Chilblains

COVID toes

Acral lesions

Serology tests

RT-PCR

Pandemic

ABSTRACT

Background: Numerous cases of chilblains have been observed, mainly in young subjects with no or mild symptoms compatible with COVID-19. The pathophysiology of these lesions is still widely debated, and an association with SARS-CoV-2 infection remains unconfirmed.

Objectives: This paper focuses on the unresolved issues about these *COVID toes* and, in particular, whether or not they are associated with COVID-19.

Arguments: The temporal link between the outbreak of chilblains and the COVID-19 pandemic suggests a link between the two events. Positive anti-SARS-CoV/SARS-CoV-2 immunostaining on skin biopsy of chilblains seems to confirm the presence of the virus in the lesions but lacks specificity and must be interpreted with caution. Conversely, RT-PCR and anti-SARS-CoV-2 serology were negative in the majority of patients with chilblains. Therefore, SARS-CoV-2 infection can be excluded, with relative certainty, even after accounting for possible lower immunization in mild/asymptomatic patients and for some differences in sensitivity/specificity between the tests used. Some authors hypothesize that chilblains could be the cutaneous expression of a strong type I interferon (IFN-I) response. High production of IFN-I is suggested to be associated with early viral control and may suppress antibody response. However, the absence of other cutaneous or extracutaneous symptoms, as observed in other interferonopathies, raises unanswered questions. To date, a direct link between chilblains and COVID-19 still seems impossible to confirm. A more indirect association due to lifestyle changes induced by lockdown is a possible explanation. Improvement of chilblains when protective measures were adopted and after lifting of lockdown supports this hypothesis.

Conclusion: Conflicting current evidence highlights the need for systematic and repeated testing of larger numbers of patients and the need for valid follow-up data that take into consideration epidemic curves and evolution of lockdown measures.

© 2020 The Author(s). Published by Elsevier Ltd on behalf of International Society for Infectious Diseases. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Skin manifestations are considered to be infrequent presentations of COVID-19, but no causal link has been formally demonstrated to date (Daneshgaran et al., 2020; Freeman et al., 2020a). A significant number of cases of chilblains have been observed, mainly in adolescents and young adults with no or mild symptoms compatible with SARS-CoV-2 infection. An association between chilblains and COVID-19 is suspected, but the pathophysiology of these lesions is still widely debated. Although numerous publications on *chilblains*, *pseudo-chilblains*, *chilblain-like lesions*, *covid-toes*, or *acral ischemic lesions* exist, an association with SARS-CoV-2 infection remains currently unconfirmed.

The prevalence of these *COVID-toes* observed in several European countries and in the United States is difficult to assess,

with some authors drawing attention to possible *SALAMI publishing* and to overlapping cases reported in scientific literature and social networks (Kluger, 2020). These lesions seem to mainly affect the feet of children, adolescents, and young adults who are otherwise in good health and who have no particular medical history (Figure 1). However, in our series, blood tests revealed isolated positive anti-nuclear antibodies in one-third of patients, and these lesions seem to preferentially affect patients with a low BMI (Median:19.13) (Baeck et al., 2020c; Herman et al., 2020). Several patients report symptoms consistent with SARS-CoV-2 infection in the days and weeks prior to the onset of chilblains or report contact with people who experienced such symptoms.

In most reported series, other causes of chilblains such as coagulopathy or systemic diseases were excluded. Several authors performed histopathological and/or immunofluorescence analyses that confirmed the diagnosis of chilblains sometimes with vasculitis and microthromboses (Herman et al., 2020; Kanitakis et al., 2020). These features are classically encountered in

* Corresponding author.

E-mail addresses: marie.baeck@uclouvain.be (M. Baeck), Anne.herman@uclouvain.be (A. Herman).



Figure 1. Clinical aspect of the COVID toes in an otherwise healthy teenager, with purplish-erythematous macules located on the toes.

chilblains and should not be confused with the ischemic acral lesions due to thrombotic vasculopathy and systemic procoagulant state observed in patients with severe or critical COVID-19 and mainly triggered by endothelial damage (probably due to direct viral effect and perivascular inflammation) (Baeck et al., 2020a; Zhang et al., 2020; Magro et al., 2020).

The discussion focuses on whether or not chilblains are associated with COVID-19:

- The temporal link between the outbreak of chilblains and the COVID-19 pandemic (moreover, during an unusual season for this type of lesion) is a first argument to suggest a link between the two events.
- Some authors have also shown positive anti-SARS-CoV/SARS-CoV-2 immunostaining on skin biopsy specimens of chilblains, which seem to confirm the presence of the virus in the lesions (Colmenero et al., 2020; Santonja et al., 2020). However, due to lack of specificity, these immunostainings should be interpreted with caution (Baeck et al., 2020b; Ko et al., 2020).
- Conversely, RT-PCR and anti-SARS-CoV-2 serology were mostly negative (Baeck et al., 2020c; Caselli et al., 2020; Colonna et al., 2020; Denina et al., 2020; El Hachem et al., 2020; Freeman et al., 2020b; Garcia-Lara et al., 2020; García-Legaz Martínez et al., 2020; Herman et al., 2020; Le Cleach et al., 2020; Lesort et al., 2020; Mahieu et al., 2020; Neri et al., 2020; Rizzoli et al., 2020; Roca-Ginés et al., 2020; Rouanet et al., 2020; Stavert et al., 2020). Negative RT-PCR on nasopharyngeal swabs could suggest that chilblains are a late symptom of COVID-19. However, serological tests were often negative. A few publications reported positive serology for anti-SARS-CoV-2 IgA (El Hachem et al., 2020; Hubiche et al., 2020). However, the sensitivity of these antibodies is questionable, and false positives arising due to excessive sensitivity cannot be ruled out (Jääskeläinen et al., 2020). Negative RT-PCR and serological tests in the majority of patients with chilblains allow the exclusion of SARS-CoV-2 infection with relative certainty, even when accounting for possible lower immunization in mild/asymptomatic patients and for some differences in the sensitivity/specificity between the tests used. For a virus-associated phenomenon, the strikingly low rate of testing positivity raises questions.
- Some authors hypothesize that chilblains could be the cutaneous expression of a strong type I interferon (IFN-I) response (Lipsker, 2020; Battesti et al., 2020; Damsky et al., 2020; Rodríguez-Villa Lario et al., 2020). A high production of IFN-I is suggested to be associated with early viral control and mild course of SARS-CoV-2 infection. IFN-I may also suppress

antibody response, which could explain why these patients would not develop antibodies and fail serologic detection. However, some questions remain unanswered with regard to the sometimes long delay between the suspected COVID-19 symptoms and the appearance of the skin lesions or with regard to the absence of other cutaneous or extracutaneous symptoms as observed in other interferonopathies (Rodero and Crow, 2016; Papa et al., 2020).

- To date, a direct link between chilblains and COVID-19 still seems impossible to confirm. Our study, like others, points towards a more indirect association between the outbreak of chilblains and the COVID-19 pandemic due to lifestyle changes induced by lockdown—in particular, in sedentary behaviours and bare-feet exposure to cool indoors (Baeck et al., 2020c; Herman et al., 2020; Neri et al., 2020; Roca-Ginés et al., 2020). Patient follow-up supports this hypothesis with the observed improvement of chilblains when protective and warm-up measures were adopted and containment restrictions were relaxed. It should also be noted that we have observed no new cases of chilblains since lifting of lockdown measures, despite recent new COVID-19 infections.

Shedding light on unanswered questions would be significant for patient management and information, testing strategies, and implementation of isolation measures. Only data on large series of patients with systematic and repeated testing, as well as follow-up information considering epidemic curves and evolution of lockdown measures, will enable to strengthen any one hypothesis.

Funding source

The authors disclose no financial associations or funding.

Ethical approval

The data have been approved by the appropriate ethics committee and have therefore been performed in accordance with the ethical standards laid down in the 1964 Declaration of Helsinki and its later amendments. Specific national laws have also been observed.

Authors' contribution

Marie Baeck and Anne Herman performed literature searches, prepared the manuscript, reviewed and approved the manuscript, and decided to submit the manuscript for publication.

Conflict of interest

The authors have no conflicts of interest.

Acknowledgments

The patients in this manuscript have given written informed consent for publication of their case details.

References

- Baeck M, Herman A, Peeters C, Marot L, Hermans C. Are chilblains a skin expression of COVID-19 microangiopathy? [published online ahead of print, 2020 July 13]. *J Thromb Haemost* 2020a;., doi:<http://dx.doi.org/10.1111/jth.15008>. 10.1111/jth.15008.
- Baeck M, Hoton D, Marot L, Herman A. Chilblains and COVID-19: why SARS-CoV-2 endothelial infection is questioned [published online ahead of print, 2020 August 15]. *Br J Dermatol* 2020b;., doi:<http://dx.doi.org/10.1111/bjd.19489>.
- Baeck M, Peeters C, Herman A. Chilblains and COVID-19: further evidence against a causal association [published online ahead of print, 2020 August 31]. *J Eur Acad Dermatol Venereol* 2020c;., doi:<http://dx.doi.org/10.1111/jdv.16901>.

- Battesti G, El Khalifa J, Abdelhedi N, Ferre V, Bouscarat F, Picard-Dahan C, et al. New insights in COVID-19-associated chilblains: a comparative study with chilblain lupus erythematosus [published online ahead of print, 2020 July 3]. *J Am Acad Dermatol* 2020;83(4):1219–22, doi:http://dx.doi.org/10.1016/j.jaad.2020.06.1018 S0190-9622(20)32147-2; Epub 2020 Jul 3.
- Caselli D, Chironna M, Loconsole D, Nigri L, Mazzotta F, Bonamonte D, et al. No evidence of SARS-CoV-2 infection by polymerase chain reaction or serology in children with pseudo-chilblain [published online ahead of print, 2020 July 1]. *Br J Dermatol* 2020;183(4):784–5, doi:http://dx.doi.org/10.1111/bjd.19349 Epub 2020 Jul 29.
- Colmenero I, Santonja C, Alonso-Riño M, Noguera-Morel L, Hernández-Martín A, Andina D, et al. SARS-CoV-2 endothelial infection causes COVID-19 chilblains: histopathological, immunohistochemical and ultrastructural study of 7 paediatric cases [published online ahead of print, 2020 June 20]. *Br J Dermatol* 2020;183(4):729–37, doi:http://dx.doi.org/10.1111/bjd.19327 Epub 2020 Aug 5.
- Colonna C, Spinelli F, Monzani NA, Ceriotti F, Gelmetti C. Chilblains in children in the time of Covid-19: new evidence with serology assay [published online ahead of print, 2020 June 13]. *Pediatr Dermatol* 2020;, doi:http://dx.doi.org/10.1111/pde.14269.
- Damsky W, Peterson D, King B. When interferon tiptoes through COVID-19: pernio-like lesions and their prognostic implications during SARS-CoV-2 infection. *J Am Acad Dermatol* 2020;83(3):e269–70, doi:http://dx.doi.org/10.1016/j.jaad.2020.06.052.
- Daneshgarian G, Dubin DP, Gould DJ. Cutaneous manifestations of COVID-19: an evidence-based review. *Am J Clin Dermatol* 2020;21(5):627–39, doi:http://dx.doi.org/10.1007/s40257-020-00558-4.
- Denina M, Pellegrino F, Morotti F, Coppo P, Bonsignori IM, Garazzino S, et al. All that glisters is not COVID: low prevalence of seroconversion against SARS-CoV-2 in a pediatric cohort of patients with Chilblain-like lesions [published online ahead of print, 2020 August 8]. *J Am Acad Dermatol* 2020;, doi:http://dx.doi.org/10.1016/j.jaad.2020.08.021 S0190-9622(20)32409-9; Epub ahead of print.
- El Hachem M, Diociaiuti A, Concato C, Carsetti R, Carnevale C, Ciofi Degli Atti M, et al. A clinical, histopathological and laboratory study of 19 consecutive Italian paediatric patients with chilblain-like lesions: lights and shadows on the relationship with COVID-19 infection [published online ahead of print, 2020 May 31]. *J Eur Acad Dermatol Venereol* 2020;, doi:http://dx.doi.org/10.1111/jdv.16682 Epub ahead of print.
- Freeman EE, McMahon DE, Lipoff JB, Rosenbach M, Kovarik C, Desai SR, et al. The spectrum of COVID-19-associated dermatologic manifestations: an international registry of 716 patients from 31 countries [published online ahead of print, 2020 July 2]. *J Am Acad Dermatol* 2020a;, doi:http://dx.doi.org/10.1016/j.jaad.2020.06.1016 S0190-9622(20)32126-5.
- Freeman EE, McMahon DE, Lipoff JB, Rosenbach M, Kovarik C, Takeshita J, et al. Pernio-like skin lesions associated with COVID-19: a case series of 318 patients from 8 countries. *J Am Acad Dermatol* 2020b;83(2):486–92, doi:http://dx.doi.org/10.1016/j.jaad.2020.05.109 Epub 2020 May 30.
- García-Lara G, Linares-González L, Ródenas-Herranz T, Ruiz-Villaverde R. Chilblain-like lesions in pediatrics dermatological outpatients during the COVID-19 outbreak [published online ahead of print, 2020 May 6]. *Dermatol Ther* 2020; e13516, doi:http://dx.doi.org/10.1111/dth.13516.
- García-Legaz Martínez M, Martínez-Doménech Á, Magdaleno-Tapiá J, Valenzuela-Oñate C, Partarrieu-Mejías F, Lorca-Spröhnle J, et al. Acute acral cutaneous manifestations during the COVID-19 pandemic: a single-centre experience [published online ahead of print, 2020 June 26]. *J Eur Acad Dermatol Venereol* 2020;, doi:http://dx.doi.org/10.1111/jdv.16777 Epub ahead of print.
- Herman A, Peeters C, Verroken A, Tromme I, Tennstedt D, Marot L, et al. Evaluation of chilblains as a manifestation of the COVID-19 pandemic [published online ahead of print 2020 June 25]. *JAMA Dermatol* 2020;e202368, doi:http://dx.doi.org/10.1001/jamadermatol.2020.2368.
- Hubiche T, Le Duff F, Chiaverini C, Giordanengo V, Passeron T. Negative SARS-CoV-2 PCR in patients with chilblain-like lesions [published online ahead of print, 2020 June 18]. *Lancet Infect Dis* 2020;, doi:http://dx.doi.org/10.1016/S1473-3099(20)30518-1 S1473-3099(20)30518-1.
- Jääskeläinen AJ, Kuivanen S, Kekäläinen E, Ahava MJ, Loginov R, Kallio-Kokko H, et al. Performance of six SARS-CoV-2 immunoassays in comparison with microneutralisation [published online ahead of print, 2020 June 15]. *J Clin Virol* 2020;129:104512, doi:http://dx.doi.org/10.1016/j.jcv.2020.104512 Epub 2020 Jun 15.
- Kanitakis J, Lesort C, Danset M, Julien D. Chilblain-like acral lesions during the covid-19 pandemic (“covid toes”): histologic, immunofluorescence and immunohistochemical study of 17 cases. *J Am Acad Dermatol* 2020;83(3):870–5, doi:http://dx.doi.org/10.1016/j.jaad.2020.05.145.
- Kluger N. Pitfalls of possible reporting of same patients with COVID-19 in dermatology journals [published online ahead of print, 2020 June 3]. *J Eur Acad Dermatol Venereol* 2020;, doi:http://dx.doi.org/10.1111/jdv.16690.
- Ko CJ, Harigopal M, Damsky W, Gehlhausen JR, Bosenberg M, Patrignelli R, et al. Perniosis during the COVID-19 pandemic: negative anti-SARS-CoV-2 immunohistochemistry in six patients and comparison to pernioles before the emergence of SARS-CoV-2 [published online ahead of print, 2020 August 3]. *J Cutan Pathol* 2020;, doi:http://dx.doi.org/10.1111/cup.13830 Epub ahead of print.
- Le Cleach L, Dousset L, Assier H, Fourati S, Barbarot S, Boulard C, et al. Most chilblains observed during the COVID-19 outbreak occur in patients who are negative for COVID-19 on PCR and serology testing [published online ahead of print, 2020 July 6]. *Br J Dermatol* 2020;183(5):866–74, doi:http://dx.doi.org/10.1111/bjd.19377 Epub 2020 Aug 9.
- Lesort C, Kanitakis J, Villani A, Ducroux E, Bouschon P, Fattouh K, et al. COVID-19 and outbreak of chilblains: are they related? [published online ahead of print, 2020 June 27]. *J Eur Acad Dermatol Venereol* 2020;, doi:http://dx.doi.org/10.1111/jdv.16779 Epub ahead of print.
- Lipsker D. A chilblain epidemic during the COVID-19 pandemic. A sign of natural resistance to SARS-CoV-2? [published online ahead of print, 2020 June 6]. *Med Hypotheses* 2020;144:109959, doi:http://dx.doi.org/10.1016/j.mehy.2020.109959.
- Magro C, Mulvey JJ, Berlin D, Nuovo G, Salvatore S, Harp J, et al. Complement associated microvascular injury and thrombosis in the pathogenesis of severe COVID-19 infection: a report of five cases. *Transl Res* 2020;220:1–13, doi:http://dx.doi.org/10.1016/j.trsl.2020.04.007 Epub 2020 Apr 15.
- Mahieu R, Tillard L, Le Guillou-Guillemette H, Vinatier E, Jeannin P, Crôuë A, et al. No antibody response in acral cutaneous manifestations associated with COVID-19? [published online ahead of print, 2020 June 2]. *J Eur Acad Dermatol Venereol* 2020;34(10):e546–8, doi:http://dx.doi.org/10.1111/jdv.16688.
- Neri I, Virdi A, Corsini I, Guglielmo A, Lazzarotto T, Gabrielli L, et al. Major cluster of pediatric “true” primary chilblains during the COVID-19 pandemic: a consequence of lifestyle changes due to lockdown [published online ahead of print, 2020 June 13]. *J Eur Acad Dermatol Venereol* 2020;, doi:http://dx.doi.org/10.1111/jdv.16751 Epub ahead of print.
- Papa R, Volpi S, Gattorno M. Type I interferonopathies: cutaneous vasculopathy, chilblains, panniculitis-induced lipodystrophy and others skin manifestations [published online ahead of print, 2020 July 2]. *G Ital Dermatol Venereol* 2020;, doi:http://dx.doi.org/10.23736/S0392-0488.20.06709-7.
- Rizzoli L, Collini L, Magnano M, Termine S, Barcelli R, Infusino SD, et al. Chilblain-like lesions during the COVID-19 pandemic: a serological study on a case series [published online ahead of print, 2020 July 1]. *Br J Dermatol* 2020;183(4):782–4, doi:http://dx.doi.org/10.1111/bjd.19348 Epub 2020 Jul 28.
- Roca-Ginés J, Torres-Navarro I, Sánchez-Arráez J, Abril-Pérez C, Sabalza-Baztán O, Pardo-Granell S, et al. Assessment of acute acral lesions in a case series of children and adolescents during the COVID-19 pandemic [published online ahead of print, 2020 June 25]. *JAMA Dermatol* 2020;156(9):992–7, doi:http://dx.doi.org/10.1001/jamadermatol.2020.2340.
- Rodero MP, Crow YJ. Type I interferon-mediated monogenic autoinflammation: the type I interferonopathies, a conceptual overview. *J Exp Med* 2016;213(12):2527–38, doi:http://dx.doi.org/10.1084/jem.20161596.
- Rodríguez-Villa Lario A, Vega-Díez D, González-Cañete M, Gómez-Zubiaur A, Pérez-Mesonero R, Bandini M, et al. Histological findings in chilblain lupus-like COVID lesions: in search of an answer to understand their aetiology [published online ahead of print, 2020 June 12]. *J Eur Acad Dermatol Venereol* 2020;34(10):e572–4, doi:http://dx.doi.org/10.1111/jdv.16733 Epub 2020 Jul 9.
- Rouanet J, Lang E, Beltzung F, Evrard B, Henquell C, Joulie I, et al. Recent outbreak of chilblain-like lesions is not directly related to SARS-CoV-2 infection [published online ahead of print, 2020 June 26]. *J Eur Acad Dermatol Venereol* 2020;, doi:http://dx.doi.org/10.1111/jdv.16776 Epub ahead of print.
- Santonja C, Heras F, Núñez L, Requena L. COVID-19 chilblain-like lesion: immunohistochemical demonstration of SARS-CoV-2 spike protein in blood vessel endothelium and sweat gland epithelium in a polymerase chain reaction-negative patient [published online ahead of print, 2020 June 25]. *Br J Dermatol* 2020;, doi:http://dx.doi.org/10.1111/bjd.19338. 10.1111/bjd.19338.
- Stavert R, Meydani-Korb A, de Leon D, Osgood R, Blau J, Luu T. Evaluation of SARS-CoV-2 antibodies in 24 patients presenting with chilblains-like lesions during the COVID-19 pandemic [published online ahead of print, 2020 August 18]. *J Am Acad Dermatol* 2020;, doi:http://dx.doi.org/10.1016/j.jaad.2020.08.049 S0190-9622(20)32440-3.
- Zhang Y, Cao W, Xiao M, Li YJ, Yang Y, Zhao J, et al. Clinical and coagulation characteristics of 7 patients with critical COVID-2019 pneumonia and acidosis. *Zhonghua Xue Ye Xue Za Zhi* 2020;41(4):302–7, doi:http://dx.doi.org/10.3760/cma.j.issn.0253-2727.2020.008.